

	FRANKSTON PRIMARY SCHOOL First Aid - Anaphylaxis Policy	Last Review 2026 Next Review: Feb 2027
---	---	--



Help for non-English speakers

If you need help to understand the information in this policy please contact the school office on 9783 3769.

PURPOSE

To explain to Frankston Primary School parents, carers, staff and students the processes and procedures in place to support students diagnosed as being at risk of suffering from anaphylaxis.

This policy also ensures that Frankston Primary School is compliant with Ministerial Order 706 and the Department's guidelines for anaphylaxis management.

SCOPE

This policy applies to:

- all staff, including casual relief staff and volunteers
- all students who have been diagnosed with anaphylaxis, or who may require emergency treatment for an anaphylactic reaction, and their parents and carers.

POLICY

Statement

Frankston Primary School will fully comply with Ministerial Order 706 and the associated guidelines published by the Department of Education and Training.

Anaphylaxis

Anaphylaxis is a severe allergic reaction that occurs after exposure to an allergen.

The most common allergens for school-aged children are nuts, eggs, cow's milk, fish, shellfish, wheat, soy, sesame, latex, certain insect stings and medication.

Symptoms

Signs and symptoms of a mild to moderate allergic reaction can include:

- swelling of the lips, face and eyes
- hives or welts
- tingling in the mouth.

Signs and symptoms of a mild to moderate allergic reaction, can include:

- swelling of the lips, face and eyes
- hives or welts
- tingling in the mouth.

Signs and symptoms of a anaphylaxis, a severe allergic reaction, can include:

- difficult/noisy breathing
- swelling of tongue
- difficulty talking and/or hoarse voice
- wheeze or persistent cough
- persistent dizziness or collapse
- student appears pale or floppy
- abdominal pain and/or vomiting.

Symptoms usually develop within ten minutes and up to two hours after exposure to an allergen but can appear within a few minutes.

Treatment

Adrenaline given as an injection into the muscle of the outer mid-thigh is the first aid treatment for anaphylaxis.

Individuals diagnosed as being at risk of anaphylaxis are prescribed an adrenaline device for use in an emergency. These adrenaline autoinjectors are designed so that anyone can use them in an emergency.

Individual Anaphylaxis Management Plans

All students at Frankston Primary School who are diagnosed by a medical practitioner as being at risk of suffering from an anaphylactic reaction must have an Individual Anaphylaxis Management Plan. When notified of an anaphylaxis diagnosis, the Principal or Principal's delegate of Frankston Primary School is responsible for developing a plan in consultation with the student's parents/carers.

Where necessary, an Individual Anaphylaxis Management Plan will be in place as soon as practicable after a student enrolls at Frankston Primary School and where possible, before the student's first day.

Parents and carers must:

- obtain an ASCIA Action Plan for Anaphylaxis (RED) from the student's medical practitioner and provide a copy to the school as soon as practicable
- immediately inform the school in writing if there is a relevant change in the student's medical condition and obtain an updated ASCIA Action Plan for Anaphylaxis (RED)

- provide an up-to-date photo of the student for the ASCIA Action Plan for Anaphylaxis (RED) when that Plan is provided to the school and each time it is reviewed
- provide the school with a current adrenaline autoinjector for the student that has not expired
- participate in annual reviews of the student's Individual Anaphylaxis Management Plan that is prepared by the school.

Each student's Individual Anaphylaxis Management Plan must include:

- information about the student's allergies and the potential for anaphylactic reaction, including the type of allergies the student has
- information about the signs or symptoms the student might exhibit in the event of an allergic reaction based on a written diagnosis from a medical practitioner
- strategies to minimise the risk of exposure to known allergens while the student is under the care or supervision of school staff, including in the school yard, at camps and excursions, or at special events conducted, organised or attended by the school
- the name of the person(s) responsible for implementing the risk minimisation strategies, which have been identified in the Individual Anaphylaxis Management Plan
- information about where the student's medication will be stored
- the student's emergency contact details
- an up-to-date ASCIA Action Plan for Anaphylaxis (RED) completed by the student's medical practitioner.

Review and updates to Individual Anaphylaxis Management Plans

A student's Individual Anaphylaxis Management Plan will be reviewed and updated on an annual basis in consultation with the student's parents/carers. The plan will also be reviewed and, where necessary, updated in the following circumstances:

- as soon as practicable after the student has an anaphylactic reaction at school
- if the student's medical condition, insofar as it relates to allergy and the potential for anaphylactic reaction, changes
- when the student is participating in an off-site activity, including camps and excursions, or at special events including fetes and concerts.

Our school may also consider updating a student's Individual Anaphylaxis Management Plan if there is an identified and significant increase in the student's potential risk of exposure to allergens at school.

Location of plans and adrenaline autoinjectors

A copy of each student's Individual Anaphylaxis Management Plan will be stored with their ASCIA Action Plan for Anaphylaxis (RED) at First Aid (sick bay), together with the student's adrenaline autoinjector (unless a student's Individual Anaphylaxis Management Plan sites different strategies). Adrenaline autoinjectors must be labelled with the student's name.

Risk Minimisation Strategies

To reduce the risk of a student suffering from an anaphylactic reaction at Frankston Primary School, we have put in place the following strategies:

Classroom	*staff and students are regularly reminded to wash their hands after eating *students are discouraged from sharing food
Yard	*students are discouraged from eating outside or taking rubbish outside *garbage bins in the yard are covered to reduce the risk of attracting insects *gloves / tongs are encouraged to be worn when picking up papers or rubbish in the playground
Canteen	*school canteen staff are trained in appropriate food handling to reduce the risk of cross-contamination
General	*a general use adrenaline autoinjector is stored in the School First Aid room *school staff complete relevant anaphylaxis training and are verified by trained school staff and complete mandated six monthly briefings *all staff informed of students at risk of anaphylaxis and requirements in student plan *planning for off-site activities will include risk minimisation strategies for students at risk of anaphylaxis including supervision requirements, appropriate number of trained staff and other risk controls appropriate to the activity and students attending * the Principal will complete the Department's Annual Risk Management Checklist for anaphylaxis management to assist with the evaluation and review of this policy and the support provided to students at risk of anaphylaxis.

Chapter 8 of the Department's [Anaphylaxis Guidelines](#) includes further detailed risk mitigation strategies that Frankston Primary School may choose to adopt dependent on the Plan and needs of each student.

Adrenaline autoinjectors for general use

Note: For general advice on the appropriate number of general use adrenaline autoinjectors, refer to chapter 10 of the Department's [Anaphylaxis Guidelines](#)

Frankston Primary School will maintain a supply of at least TWO adrenaline autoinjector(s) for general use, as a back-up to those provided by parents and carers for specific students, and also for students who may suffer from a first-time reaction at school.

There are currently 4 adrenaline devices approved by the Therapeutic Goods Administration for use in Australia:

- the EpiPen®
- the Anapen®
- Jext®
- Neffy®.

All devices can be used when provided by families for students, however, the Principal or allocated staff member can only use EpiPen®, Anapen® or Jext® adrenaline autoinjector for general use. For more information about which autoinjector to purchase for general use, refer to [Adrenaline autoinjectors for general use](#).

Adrenaline autoinjectors for general are stored at First Aid (sick bay) in the main school office and labelled "general use".

The Principal's delegate is responsible for arranging the purchase of adrenaline autoinjectors for general use, and considers:

- the number of students enrolled at Frankston Primary School at risk of anaphylaxis
- the accessibility of adrenaline autoinjectors supplied by parents/carers
- the availability of a sufficient supply of autoinjectors for general use in different locations at the school, as well as at camps, excursions and events
- the limited life span of adrenaline autoinjectors, and the need for general use adrenaline autoinjectors to be replaced when used or prior to expiry
- the weight of the students at risk of anaphylaxis to determine the correct dosage of adrenaline autoinjector/s to purchase.

Emergency Response

In the event of an anaphylactic reaction, the emergency response procedures in this policy must be followed, together with the school's general first aid procedures, emergency response procedures and the student's Individual Anaphylaxis Management Plan.

A complete and up-to-date list of students identified as being at risk of anaphylaxis is maintained by the School First Aid officer and stored at First Aid. For camps, excursions and special events, a designated staff member will be responsible for maintaining a list of students at risk of anaphylaxis attending the special event, together with their Individual Anaphylaxis Management Plans and adrenaline devices, where appropriate.

If a student experiences an anaphylactic reaction at school or during a school activity, school staff must:

Step	Action
1.	Lay the person flat Do not allow them to stand or walk If breathing is difficult, allow them to sit with legs outstretched Be calm and reassuring Do not leave them alone Seek assistance from another staff member or reliable student to locate the student's adrenaline device or the school's general use autoinjector, and the student's Individual Anaphylaxis Management Plan, stored at First Aid If the student's plan is not immediately available, or they appear to be experiencing a first time reaction, follow steps 2 to 5
2.	Administer an EpiPen or EpiPen Jr Remove from plastic container Form a fist around the EpiPen and pull off the blue safety release (cap) Hold leg still and place orange end against the student's outer mid-thigh (with or without clothing) Push down hard until a click is heard or felt and hold in place for 3 seconds Remove EpiPen Note the time the EpiPen is administered Retain the used EpiPen to be handed to ambulance paramedics along with the time of administration. OR

	Administer an Anapen® 500 Pull off the black needle shield Pull off grey safety cap (from the red button) Place needle end firmly against the student's outer mid-thigh at 90 degrees (with or without clothing) Press red button so it clicks and hold for 3 seconds Remove Anapen® Note the time the Anapen is administered Retain the used Anapen to be handed to ambulance paramedics along with the time of administration. OR Administer Jext 150 or 300 Form fist around Jext and pull off yellow cap Place black injector tip against outer-mid thigh (with or without clothing) Push black tip firmly until a click is heard and hold in place for 3 seconds. Remove Jext Note the time the Jext device is administered. The used adrenaline device must be handed to the ambulance paramedics along with the time of administration OR Administer Neffy® 1mg or 2mg Hold the nasal spray with your thumb on the bottom of the plunger and a finger on either side of the nozzle. Do not pull or push on the plunger. Do not test or prime (pre-spray). Each Neffy nasal spray contains only one spray. Place the nozzle of the nasal spray into a nostril until fingers touch the nose. For smaller nostrils, aim for the fingers to touch the nose. Keep the nozzle pointed towards the forehead. Do not angle the nozzle of the nasal spray to the inner or outer walls of the nose. Press the plunger up firmly until the dose is administered and it sprays into the nostril. Note the time the Neffy device is administered. The used adrenaline device must be handed to the ambulance paramedics along with the time of administration
--	---

3.	Call an ambulance (000)
4.	If there is no improvement or severe symptoms progress (as described in the ASCIA Action Plan for Anaphylaxis (RED)), further adrenaline doses may be administered every 5 minutes, if other adrenaline autoinjectors are available.
5.	Contact the student's emergency contacts.
6.	The Principal or Principal delegate, must contact the Incident Support and Operations Centre (ISOC) on 1800 126 126 to report 'High' or 'Extreme' severity incidents to report the incident. Incidents assessed as 'Low' or 'Medium' can be reported directly into EduSafe Plus by the Principal or allocated staff member.

If a student appears to be having a severe allergic reaction, but has not been previously diagnosed with an allergy or being at risk of anaphylaxis, school staff should follow steps 2 – 5 as above.

For first time anaphylactic reactions, the school's general use adrenaline autoinjector device must be used. If the general use device is not immediately available in an anaphylaxis emergency, staff may use another student's adrenaline device, including the Epipen®, Anapen®, Jext® or Neffy® device. This may save a life.

If another student's adrenaline device is used in an anaphylaxis emergency, the school must notify the parents of the student whose device was used and immediately replace the device.

Where possible, schools should consider using the correctly dosed adrenaline device depending on the weight of the student. However, in an emergency if there is no other option available, any device should be administered to the student.

COMMUNICATION PLAN

This policy will be available on Frankston Primary School's website so that parents and other members of the school community can easily access information about Frankston Primary School's anaphylaxis management procedures.

The Principal or Principal's delegate is responsible for ensuring that all relevant staff, including casual relief staff and canteen staff are aware of this policy and Frankston Primary School's procedures for anaphylaxis management.

Casual relief staff who are responsible for the care and/or supervision of students who are identified as being at risk of anaphylaxis will receive a verbal briefing on this policy, their role in responding to an anaphylactic reaction and where required, the identity of students at risk.

We will also communicate this policy through:

- Staff briefings
- School Newsletter

Staff training

The Principal is responsible for ensuring relevant staff are trained and briefed in anaphylaxis management, consistent with the department's [Anaphylaxis Guidelines](#). Including:

- School staff who conduct classes attended by students who are at risk of anaphylaxis

Staff who are required to undertake training must have completed:

- an approved face-to-face anaphylaxis management training course in the last three years, or
- an approved online anaphylaxis management training course in the last two years.

Frankston Primary School uses the following department recommended ASCIA eTraining course for Victorian Schools with 22579VIC, or 22578VIC followed by a competency check by the school anaphylaxis supervisor.

Staff are also required to attend a briefing on anaphylaxis management and this policy at least twice per year (with the first briefing to be held at the beginning of the school year), facilitated by a staff member who has successfully completed an anaphylaxis management course within the last 2 years including our School Anaphylaxis Supervisor. Each briefing will address:

- the school's anaphylaxis management policy
- the causes, symptoms and treatment of anaphylaxis
- the identities of students diagnosed as being at risk of anaphylaxis, their allergens and the location of their Individual Anaphylaxis Management Plans and their medication/s
- discussion on staff anaphylaxis training and renewal
- how to use an adrenaline device, including hands-on practice with an adrenaline device trainer device (which does not contain adrenaline)
- the school's general first aid and emergency procedures
- the location of adrenaline devices prescribed for individual students that have been purchased by their family
- how to access on-going support and training.

When a new student enrolls at Frankston Primary School who is at risk of anaphylaxis, the Principal or Principal's delegate will develop an interim plan in consultation with the student's parents and ensure that appropriate staff are trained and briefed as soon as possible.

A record of staff training courses and briefings will be maintained through the Staff Training Register – reflecting the mandate of twice-yearly briefing sessions and evidence of compliance with the training requirements of Ministerial Order 706 – Anaphylaxis Management in Victorian Schools. The record will include the names of staff who have undertaken the training course and the date the training is due for renewal, as well as the names of the staff who attended the twice yearly briefing to staff.

The Principal will ensure that while students at risk of anaphylaxis are under the care or supervision of the school outside of normal class activities, including in the school yard, at camps and excursions, or at special event days, there is a sufficient number of school staff present who have been trained in anaphylaxis management.

Communication

This policy will be communicated to our school community in the following ways:

- Available publicly on our school's website
- Reminders in our school newsletter
- Hard copy available from school administration upon request

FURTHER INFORMATION AND RESOURCES

- The department's Policy and Advisory Library (PAL):
- [Anaphylaxis](#)
- [Allergies](#)
- [First Aid for Students and Staff](#)
- [Health Care Needs](#)
- Managing Reporting School Incidents (Including Emergencies)
- [Medication](#)
- [Allergy & Anaphylaxis Australia](#)
- ASCIA Guidelines: [Schooling and childcare](#)
- [Hero HQ Anaphylaxis Management Training](#)
- <https://allergyfacts.org.au/interest/anaphylaxis/>
- Royal Children's Hospital: [Allergy and immunology](#)
- Frankston Primary School's First Aid policies on [school website](#)

NEXT REVIEW	February 2027
PRINCIPAL	Renée Kennedy
EVALUATION	This policy will be reviewed annually as part of the school's review cycle, or as required due to changes in relevant Acts, Laws, DE mandates, or should situations arise that require earlier consideration.

The principal will complete the department's Annual Risk Management Checklist for anaphylaxis management to assist with the evaluation and review of this policy and the support provided to students at risk of anaphylaxis.